



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

MICROSCOPY.¹

Methods of Preparing Molluscan Ova.²—The results of my work for the first three months were not promising and it was not until I had hit upon my present methods of preparing surface views of the entire ova that any detailed study of the cleavage could be made. Since I owe most of my results to this method and since I am convinced that it may be profitably employed in the preparation of the surface views of many different objects I believe it merits a detailed description.

The ova were fixed in many different fluids—Kleinenberg's Picro-sulphuric, Picric acid in sea water, Merkel's, Perenyi's, Flemming, stronger and weaker, Auerbach's, Corrosive sublimate, Chromo-formic, Chromo-acetic and absolute alcohol—but none of these methods for a moment compare with the first named, i. e. Kleinenberg's stronger picro-sulphuric. The ova were left in this for a length of time varying from fifteen minutes to one hour and were then gradually transferred to 70% alcohol. They were left in this until all traces of picric acid had been washed out and were finally preserved in 95% alcohol. During the first year of the work many of the preparations were ruined by becoming very dark, owing I think to the extraction of tannin from the corks. This trouble was afterward avoided by using rubber corks, or better still by coating ordinary corks with a thin layer of paraffin.

As a result of many experiments with almost every one of the common staining fluids, I found that the best method of preparing surface views of the whole egg or embryo was the following:—(1) Transfer the object gradually from alcohol to water. (2) Stain from five to ten minutes in a solution of Delafield's (Grenacher's) Hæmatoxylin diluted about six times with distilled water and rendered *slightly* acid by a trace of HCl. (3) Dehydrate and clear in oil of cedar or cloves. (4) Mount in Balsam supporting the cover glass so as to prevent crushing. By occasionally softening the balsam with a drop or two of xylol and slightly moving the cover glass the objects can be rolled into any position desired.

By this method wonderfully beautiful surface preparations were obtained showing with remarkable clearness not only the nuclei and cell boundaries but also the caryokinetic figures and in many cases the archoplasmic spheres and centrosomes. One very considerable advan-

¹Edited by C. O. Whitman, Chicago University.

(²Extracted from a paper to be published later on the development.)

tage of this method is that the preparations were permanent—in fact they become better with age instead of degenerating. All the preparations from which the figures were drawn are still in existence and can be consulted at any time.

I have employed this method with almost as good results in the preparation of surface views of the embryo chick and English sparrow and also, with considerable success on other molluscan eggs and embryos as well as those of annelids and echinoderms.

—E. G. CONKLIN, Delaware, Ohio.